

# TARSAL TUNNEL SYNDROME



American College of  
Foot and Ankle Surgeons


## What Is the Tarsal Tunnel?

The tarsal tunnel is a narrow space that lies on the inside of the ankle next to the ankle bones. The tunnel is covered with a thick ligament (the flexor retinaculum) that protects and maintains the structures contained within the tunnel—arteries, veins, tendons, and nerves. One of these structures is the posterior tibial nerve, which is the focus of tarsal tunnel syndrome.

## What Is Tarsal Tunnel Syndrome?

Tarsal tunnel syndrome is a compression, or squeezing, on the posterior tibial nerve that produces symptoms anywhere along the path of the nerve. The posterior tibial nerve runs along the inside of the ankle into the foot.

Tarsal tunnel syndrome is similar to carpal tunnel syndrome, which occurs in the wrist. Both disorders arise from the compression of a nerve in a confined space. Although tarsal tunnel syndrome may not be as well known as carpal tunnel syndrome, it is nevertheless a cause of foot and ankle pain in adults.



## Symptoms


Patients with tarsal tunnel syndrome experience one or more of the following symptoms:

- Tingling, burning, or a sensation similar to an electrical shock
- Numbness
- Pain, including shooting pain

The symptoms are typically felt on the inside of the ankle and/or on the bottom of the foot. In some people, a symptom may be isolated and occur in just one spot. In others, it may extend to the heel, arch, toes, and even the calf.

Sometimes the symptoms of the syndrome appear suddenly. Often they are brought on or aggravated by overuse of the foot—such as in prolonged standing, walking, exercising, or beginning a new exercise program.

It is very important to seek early treatment if any of the symptoms of tarsal tunnel syndrome occur. If left untreated, the condition progresses and may result in permanent nerve damage. In addition, because the



Ankle cross section

symptoms of tarsal tunnel syndrome can be confused with other conditions, proper evaluation is essential so that a correct diagnosis can be made.

## Causes

Tarsal tunnel syndrome is caused by anything that produces compression on the posterior tibial nerve, such as:

- A person with flat feet is at risk for developing tarsal tunnel syndrome, because the outward tilting of the heel that occurs with “fallen” arches can produce strain and compression on the nerve.
- An enlarged or abnormal structure that occupies space within the tunnel can compress the nerve. Some examples include a varicose vein, ganglion cyst, swollen tendon, and arthritic bone spur.
- An injury, such as an ankle sprain, may produce inflammation and swelling in or near the tunnel, resulting in compression of the nerve.
- A person who is overweight may be prone to experiencing pressure on the posterior tibial nerve.
- Systemic diseases such as diabetes or arthritis can cause swelling, thus compressing the nerve.

## Diagnosis

The foot and ankle surgeon will examine the foot to arrive at a diagnosis and determine if there is any loss of feeling. During this examination, the

surgeon will position the foot and tap on the nerve to see if the symptoms can be reproduced. He or she will also press on the area to help determine if a small mass is present.

Sometimes an MRI is ordered, usually if a mass is suspected or in cases where initial treatment does not reduce the symptoms. In addition, special studies used to evaluate nerve problems—electromyography and nerve conduction velocity (EMG/NCV)—may be ordered if the condition shows no improvement with non-surgical treatment.

## Treatment

A variety of treatment options, often used in combination, are available to treat tarsal tunnel syndrome.

These include:

- **Rest.** Staying off the foot prevents further injury and

encourages healing.

- **Ice.** To reduce swelling in the tarsal tunnel, apply a bag of ice over a thin towel to the affected area for 20 minutes of each waking hour. Do not put ice directly against the skin.
- **Oral medications.** Nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, help reduce the pain and inflammation.
- **Immobilization.** Restricting movement of the foot by wearing a cast is sometimes necessary to enable the nerve and surrounding tissue to heal.
- **Physical therapy.** Ultrasound therapy, exercises, and other forms of physical therapy may be prescribed to reduce symptoms.
- **Injection therapy.** Injections of a local anesthetic provide pain relief, and an injected corticosteroid may be useful in treating the inflammation.

teroid may be useful in treating the inflammation.

- **Orthotic devices.** Custom shoe inserts may be prescribed to help maintain the arch and limit excessive motion that can cause compression on the nerve.
- **Shoes.** Supportive shoes, as recommended by your foot and ankle surgeon, may prove helpful.
- **Bracing.** Patients with flatfoot or those with severe symptoms and nerve damage may be fitted with a brace to reduce the amount of pressure on the foot.
- **Surgery.** Sometimes surgery is the best option for treating tarsal tunnel syndrome. The foot and ankle surgeon will determine if surgery is necessary and will select the appropriate procedure or procedures based on the cause of the condition. ▲



Melissa J. Lockwood, D.P.M.

Melissa J. Lockwood, D.P.M. — As founder of Heartland Foot and Ankle Associates, P.C., Dr. Melissa J. Lockwood happily returns to central Illinois after seven years in Cleveland, Ohio. She is a graduate of Providence Catholic High School in New Lenox, Illinois. Dr. Lockwood completed her undergraduate degree from the University of Illinois (Urbana/Champaign) in 2001 with a Bachelor of Science in Biology and a minor degree in Chemistry. Dr. Lockwood has a great interest in archaeology and also received a minor degree in Anthropology from the U of I in 2001.

Dr. Lockwood went on to the Ohio College of Podiatric Medicine and matriculated with her Podiatric Medical Degree in 2005. While at OCPM, Dr. Lockwood served as both a delegate to the student branch of the APMA

as well as President of the Student Government organization (OPMSA). Dr. Lockwood was the recipient of several awards while attending OCPM, including the Ohio Podiatric Medical Association Dr. James Conforti Community Relations Award and the OCPM Mildred Kaufman Memorial Award for Proficiency in Orthopedics and Biomechanics.

Dr. Lockwood arrives in Bloomington following a three year residency in Cleveland, Ohio. Dr. Lockwood completed a 36 month Podiatric Medical and Surgical Residency Program based out of the University Hospital System/Richmond Heights Medical Center. Dr. Lockwood has extensive training in both forefoot (bunions, hammertoes, neuromas) elective surgical procedures as well as rearfoot reconstruction. She has a special interest in the treatment of chronic wounds as well as diabetic palliative care. Dr. Lockwood enjoys working with patients one on one in a clinical setting and chatting about her favorite sports teams!

Dr. Lockwood and her husband Scott are pleased to be back in their home state and rooting on the Illini, Redbirds, Chicago Bears, and White Sox! Scott is a 2000 Illinois State University Alumni with a BS degree in Accountancy. He is a CPA working in Bloomington.

### EDUCATION

University of Illinois, Urbana-Champaign, BS Biology (01)  
Professional: Ohio College of Podiatric Medicine, (05)  
Residency: University Hospital Richmond Medical Center (05-June 08)

### PROFESSIONAL ACCREDITATION

NBPME Part I (03) NBPME Boards Part II & Part III - PM Lexus (05)  
State of Illinois Licensure (08)

### HONORS AND AWARDS

Illinois State Scholar (1997)  
OCPM Academic Scholarship (01) OCPM Dean's List (04)  
OPMA Dr. James Conforti Community Relations Award (05)  
OCPM Mildred Kaufman Memorial Award for Proficiency in Orthopedics and Biomechanics (05)  
University Hospitals Richmond Medical Center Outstanding Podiatric Resident (08)

### PROFESSIONAL ORGANIZATIONS

President, Ohio Podiatric Medical Student's Association (02-04)  
Delegate, American Podiatric Medical Student's Association (02-04)  
Chairman, President's Committee, American Podiatric Medical Student's Association Feb. 2004  
Member - OCPM Board of Trustees (02-04)  
APMA/IPMA, AAPPM, AAWP member at large

### RESEARCH/PUBLICATIONS

*Sesamoid Pathology*, coauthored with  
Dr. Joseph Favazzo, D.P.M., December 2005. Compilation of forefoot pathology edited by Vincent J. Hetherington, D.P.M.



HEARTLAND FOOT AND ANKLE ASSOCIATES, P.C.

10 Heartland Drive, Suite B  
Bloomington, IL 61704  
Phone: 309-661-9975  
Fax: 309-661-9920  
www.heartlandfootandankle.com  
drlockwood@heartlandfootandankle.com



American College of  
Foot and Ankle Surgeons

This information has been prepared by the Consumer Education Committee of the American College of Foot and Ankle Surgeons, a professional society of 5,800 foot and ankle surgeons. Members of the College are Doctors of Podiatric Medicine who have received additional training through surgical residency programs.

The mission of the College is to promote superior care of foot and ankle surgical patients through education, research and the promotion of the highest professional standards.

Copyright © 2005, American College of Foot and Ankle Surgeons • www.FootPhysicians.com